

Confinement vs. screening in 3d (effective high T) gauge theories

Owe Philipsen

Institut für Theoretische Physik, Philosophenweg 16, D-69120 Heidelberg, Germany

The physics of electric and magnetic screening in non-Abelian finite temperature field theory is closely related to the non-perturbative properties of the corresponding three-dimensional effective theories which are confining. I discuss recent lattice results obtained for the mass spectrum, string tension and the static potential in SU(2) pure gauge theory and the SU(2) Higgs model and their connection to screening masses. The non-perturbative $O(g_3^2)$ corrections to the Debye mass are evaluated.

I. INTRODUCTION

Three-dimensional gauge theories play an important role for high temperature particle physics, since they constitute the Matsubara zero mode sector of finite temperature quantum field theories in the imaginary time formalism. In particular, if the temperatures of interest are larger than the mass scales of the theory, fermions and non-zero modes may be integrated out by dimensional reduction [1] to leave a three-dimensional effective theory describing all static properties and the equilibrium thermodynamics of the theory under consideration. While dimensional reduction is a perturbative procedure, the resulting three-dimensional gauge theories require non-perturbative techniques in order to calculate physical quantities. In particular, in the cases of QCD and the symmetric phase of the electroweak sector of the Standard Model, the three-dimensional effective theories are confining and thus not accessible by perturbative methods. It is then expedient to split the problem into two parts: perform dimensional reduction perturbatively, and study the resulting three-dimensional theory on the lattice. This approach has been particularly successful in clarifying the order and strength of the electroweak phase transition [2], and is also applied to the more difficult case of QCD and the quark gluon plasma, e.g. [3]. An important question connected with non-perturbative physics concerns the screening properties of non-Abelian plasmas like Debye screening, and the much less understood question of magnetic screening and a corresponding “magnetic mass” [4].

In this contribution, the concept of screening masses is considered in the framework of three-dimensional gauge theories. Starting point is the argument that any static physical quantity of the finite temperature field theory must have a corresponding quantity in the three-dimensional theory, with which it agrees up to a (perturbative) part due to the non-zero Matsubara modes. In general, a static screening length in finite temperature field theory is defined as the exponential decay of

some *spatial* correlation function. In the dimensionally reduced theory, the direction of the correlation may be taken to correspond to (euclidean) time, and hence the same quantity appears in the spectrum or some other physical property of the (2+1) dimensional theory. In order to fully understand screening in non-Abelian plasmas, it is therefore necessary to understand the non-perturbative physics of the underlying three-dimensional theories. Here I shall discuss the physical properties of (2+1) dimensional gauge theories like the mass spectrum, the static potential, the string tension, screening of the static potential and the various correlation functions that serve to study them. The connection of these quantities to the magnetic mass and the Debye mass is discussed. Special emphasis is put on the non-perturbative, confining nature of the theories. Correspondingly, most of the presented results are drawn from lattice simulations.

II. THE SU(2) HIGGS MODEL

As a prominent example corresponding to the dimensionally reduced electroweak Standard Model [5], consider the SU(2) Higgs model in (2+1) dimensions,

$$S = \int d^3x \operatorname{Tr} \left(\frac{1}{2} F_{ij} F_{ij} + (D_i \Phi)^\dagger D_i \Phi + m_3^2 \Phi^\dagger \Phi + 2\lambda_3 (\Phi^\dagger \Phi)^2 \right), \quad (1)$$

where all fields are in a 2×2 matrix notation. The gauge coupling g_3 and the scalar coupling λ_3 have mass dimension 1/2 and 1, respectively, so the physical properties of the theory are determined by two dimensionless parameters, $x = \lambda_3/g_3^2$ and $y = m_3^2/g_3^4$. In the framework of dimensional reduction, these parameters are of course determined by the four-dimensional couplings g, λ and temperature [5]. In the present context the interest is merely in three-dimensional dynamics and this dependence will be suppressed.

The phase diagram of the model is schematically shown in Fig. 1. The perturbative Higgs phase and the non-perturbative symmetric or confinement phase are for small x separated by a line of first order phase transitions. With increasing x the transition becomes weaker until it ends in a point with a second order phase transition [6]. Beyond this point the transition disappears entirely to become a smooth crossover, so the phases are analytically connected. This phase diagram was only established non-perturbatively in the last few years [7,8] and answered many relevant questions concerning the thermodynamics of the electroweak phase transition. Here we are mainly




FIG. 1. Schematic phase diagram of the SU(2) Higgs model with a line of first order phase transitions.

concerned with the non-perturbative properties of the confinement phase. Due to the connected nature of the phase diagram, it is possible to smoothly vary the parameters from some point in the Higgs region, where the physics is perturbative and well understood, to the confinement regime and also pure gauge theory, the latter by making the scalars infinitely heavy so that they decouple.

III. MASS SPECTRUM

All physical properties of the theory are encoded in gauge-invariant n-point functions. In particular, the mass spectrum is computed from the exponential fall-off of two-point correlation functions

$$\lim_{|x-y| \rightarrow \infty} \langle \phi^\dagger(x) \phi(y) \rangle \sim e^{-M|x-y|}, \quad (2)$$

where ϕ generically denotes some gauge-invariant operator with quantum numbers J^{PC} . For example, operators with quantum number 0^{++} are

$$\begin{aligned} R &\sim \text{Tr}(\Phi^\dagger(x)\Phi(x)), \\ L &\sim \text{Tr}((D_i\Phi)^\dagger(x)D_i\Phi(x)), \\ P &\sim \text{Tr}(F_{ij}^a(x)F_{ij}^a(x)), \end{aligned} \quad (3)$$

where R and L are typically used to measure e.g. the Higgs mass, whereas P serves to compute glueball masses in pure gauge theory. From each of these operator types analogues with quantum numbers other than 0^{++} can be constructed by taking suitable linear combinations with the desired rotational and PC -symmetry properties. A gauge-invariant 1^{--} operator, used to compute the mass of the W-boson, is given by




FIG. 2. Mass spectrum of the SU(2) Higgs model for $x = \lambda_3/g_3^2 = 0.0239$ and two values of $y = m_3^2/g_3^4$, located in the Higgs (right) and confinement (left) regions.

$$V_i^a \sim \text{Tr}(\sigma^a \Phi^\dagger(x) D_i \Phi(x)). \quad (4)$$

A method to compute the mass eigenstates including the spectrum of higher excitations is to measure the full correlation matrix between all available operators in a given quantum number channel, $\langle \phi_i^\dagger(x) \phi_j(y) \rangle$, where $\phi_i \in \{R, L, P\}$, and to diagonalise it by a variational calculation to find the mass eigenstates of the Hamiltonian (for details see [9] and references therein),

$$\varphi_i = \sum_k a_{ik} \phi_k. \quad (5)$$

The coefficients a_{ik} are a measure for the overlap or wavefunction between a given mass eigenstate φ_i and the original operators ϕ_k .

The results of a lattice calculation [9] of the lowest states of the spectrum with quantum numbers 0^{++} , 2^{++} and 1^{--} at some points in the Higgs and confinement region of the phase diagram are displayed in Fig.2. In the Higgs region we see the familiar and perturbatively calculable Higgs and W-boson in the 0^{++} and 1^{--} channels, respectively. There is a large gap to the higher excitations which are scattering states. In the confinement region, in contrast, there is a dense spectrum of bound states in all channels, very much resembling the situation in QCD. Open symbols denote bound states of scalar fields, whereas full symbols represent the low lying glueballs. This identification is based on a detailed mixing analysis employing the overlaps (3), as shown in Fig. 3 for the 0^{++} channel. There it is easy to see that the ground state is a bound state of scalar fields, with almost exclusive contributions from R, L -type operators. The first excitation is a mixture of scalar and gluonic contributions. The second excited state, in contrast, has almost exclusively gluonic content and hence corresponds to a glueball.

Furthermore, the numerical values of the glueball masses agree within less than 5% with those obtained




FIG. 3. The coefficients a_{ik} , eq. (5), of the operators eq. (3) for the lowest three 0^{++} states in the confinement region (ground state at the bottom). Indices on the operators refer to different smearing levels [9].

in lattice simulations of pure SU(2) gauge theory [10]. In the latter, the physical spectrum consists of glueballs only. It is very interesting to note that adding scalar matter fields to the pure gauge theory leaves the glueball spectrum practically undisturbed, but simply adds an apparently disjoint sector of mesonic states to the spectrum. This observation is further corroborated by the fact that the string tension in the confinement region of the Higgs model is measured to be about 97% of the one in pure gauge theory [9,11]. It has further been established in [9] that the glueball masses as well as the string tension in the confinement phase are insensitive to variations of the parameters x, y , which only influence the masses of the bound states of scalars. There are preliminary results indicating that this behaviour is repeated in the SU(2) Higgs model with scalar fields in the adjoint representation [12].

In summary, the mass spectrum in three-dimensional, non-Abelian gauge Higgs models consists of a copy of the glueball spectrum of the corresponding pure gauge theory and additional bound states of matter fields.

IV. MAGNETIC MASS VS. PHYSICAL MASS SPECTRUM

A motivation for the occurrence of the magnetic mass in hot non-Abelian gauge theories has been given in the contribution of Nair [13]. In the following sections the question is considered, whether the magnetic mass can be identified non-perturbatively in a three-dimensional gauge theory, and how it might be related to physical quantities of the theory.

In the three-dimensional theory, the magnetic mass according to its standard definition appears as a pole in the gluon propagator,

	ref.	m_A/g_3^2
1-loop gap eq.	[18]	0.38
	[7,19]	0.28
	[20]	0.25
2-loop gap eq.	[21]	0.34
lattice (Landau gauge)	[14]	0.35(1)
	[15]	0.46(3)

TABLE I. Comparison of gluon propagator masses from gap equations and lattice simulations.

$$\langle A_i^a(x) A_j^b(y) \rangle \sim e^{-m_A |x-y|}. \quad (6)$$

This definition has a caveat. The pole of the gluon propagator can be shown to be gauge-invariant order by order in (resummed) perturbation theory [16], but this does not guarantee that the full non-perturbative propagator has a pole. On the other hand, since the gluon propagator itself is gauge-variant, a fully non-perturbative lattice calculation requires gauge fixing. Since that proceeds numerically, gauge-invariance of the mass is not explicit. Furthermore, due to the nature of gauge fixing procedures on the lattice, it is also not immediately evident whether the mass signal extracted really corresponds to a pole or some other analytic structure. However, it is important to note that it is quite possible for a pole in the full gluon propagator to exist, without contradicting the fact that no asymptotic gluon states can be observed [17]. A physical asymptotic state is indicated by a pole in the correlation function of a gauge-invariant operator, and clearly the gluon propagator does not belong to this class.

The correlator (6) has been evaluated for pure gauge theory by means of gap equations in various different resummation schemes at the one loop level [7,18,19], as discussed in [13] and summarised in Table I. The difference in the results is rather small given that this is the leading order result for a fully non-perturbative quantity. A recent two-loop calculation [21] employing the resummation of [7] finds corrections of about 15% and thus one may hope for convergence of the resummed perturbation series. There are also lattice simulations of the gluon propagator in Landau gauge, see Table I. In the simulations in [15] different gauges were employed and no notable gauge dependence was found. At present, the reliability and accuracy of the calculations has not yet reached a fully satisfactory quantitative level. But there is evidence that the gluon propagator in three dimensions can be evaluated by non-perturbative methods in a fixed gauge, and that it produces a mass scale of the order of $m_A \sim 0.35 - 0.46 g_3^2$.

Given the difficulties with the propagator pole the question arises whether the magnetic mass can be identified in another manifestly gauge-invariant and non-perturbative way. If a pole exists in the full gluon propagator, does it show up in the spectrum or another phys-

ical property of the theory? Consider again the SU(2) Higgs model discussed above. In the Higgs phase, there are free point like Higgs and W-bosons described by the gauge-invariant operators R and V . The masses extracted from the correlation functions of these operators agree perfectly with those extracted from the poles of the propagators [14], as is expected in a perturbative Higgs regime. As one moves around the phase diagram to the confinement phase, the gauge-invariant correlation functions and the propagators split up. In particular, m_A extracted from the gluon propagator remains constant, i.e. it is independent of the values of the scalar parameters x, y (as it should if it represents a property of pure gauge theory), whereas the mass of the 1^{--} state rises with x and y to become much heavier than m_A , c.f. Fig. 2. This is in accord with its interpretation as a bound state of scalars. Thus the operator V is not suitable to describe the magnetic mass in the symmetric phase of the Higgs model, in pure gauge theory without scalars it cannot even be defined. Likewise it is apparent from Fig. 2 that m_A cannot be identified with any glueball mass (those with quantum numbers other than $0^{++}, 2^{++}$ are even heavier [10]). In other words, if there is a mass scale associated with the propagator, it does not show up directly in the spectrum.

V. A CONSTITUENT PICTURE

An indirect way of identifying the propagator mass in the spectrum by means of a constituent picture has been proposed in [22]. In the constituent picture, the gauge-invariant composite operators describe the asymptotic mass eigenstates, whereas the propagator masses are identified as “constituent” masses. In the Higgs phase of the SU(2) Higgs model the asymptotic states correspond to point particles, i.e. the Higgs and W-bosons exist as free particles, and hence propagators and gauge-invariant operators give an equivalent description. Moving into the confinement phase, the coupling becomes strong and what were free particles now become constituents to form bound states. The latter are the asymptotic states with poles in the correlation functions of the composite gauge-invariant operators, whereas the propagator poles give the constituent masses which do not correspond to asymptotic states. This is a unified description where only gauge-invariant operators correspond to asymptotic states. The fact that asymptotic states in the Higgs phase can also be described by the gauge-variant propagators is a consequence of the perturbative Higgs nature of that regime [22].

To the extent that such a constituent picture holds, the bound state masses should be the sum of the constituent masses plus some binding energy. Counting Φ as a constituent scalar and D_μ as a constituent vector particle, the naive mass formulae for the operators in eqs. (3), (4) are,

	$J^{PC} = 0^{++}$			$J^{PC} = 1^{--}$
	R	L	P	V
lattice [9]	0.839(15)	1.47(4)	1.60(4)	1.27(6)
const. model	-	1.76	1.67	1.29

TABLE II. Comparison of screening masses from lattice simulations and a constituent model using $m_A/g_3^2 = 0.46$, m_R is used to fix the constituent scalar mass.

$$\begin{aligned} m_R &\simeq 2m_\Phi & m_L &\simeq 2m_\Phi + 2m_A \\ m_P &\simeq 4m_A & m_V &\simeq 2m_\Phi + m_A. \end{aligned} \quad (7)$$

A comparison with the numerical values from lattice simulations is given in Table II. For a test of the constituent picture in the SU(2) adjoint Higgs model, see [15].

Clearly, this naive counting rule works only for operators that have very good overlap with a definite mass eigenstate. This is the case for R, P, V but not for L , c.f. Fig. 3. Since all binding effects have been neglected, such a constituent picture can at best be qualitatively correct. Moreover, it is not suitable to describe higher excitations. Its purpose in the present context is not to give precise predictions for the mass eigenstates, but to furnish a possible connection between propagator masses and the spectrum of gauge-invariant operators.

VI. STATIC POTENTIAL AND SCREENING

Another interesting quantity determining the physical properties of confining theories is the potential energy of static colour sources, which is calculated from the exponential decay of large Wilson loops,

$$V(r) = - \lim_{t \rightarrow \infty} \ln \left[\frac{W(r, t)}{W(r, t-1)} \right]. \quad (8)$$

As in four dimensions, a string of colour flux binds the sources (in fundamental as well as adjoint representation) causing a linear rise in the potential with growing separation r . For sources in the fundamental representation, this linear rise continues to infinity in pure gauge theory. If fundamental matter fields are present as in the Higgs model, the string breaks at some scale r_b , when its energy is large enough to produce a pair of scalars. The dynamical scalars are then bound to the static sources forming a pair of static-light “mesons”, and the Wilson loop obeys a perimeter law $W(r, t) \sim \exp[-2M(r+t)]$ [23] resulting in a saturation of the potential at a constant value corresponding to twice the static-light meson energy,

$$V(R \rightarrow \infty) = 2M. \quad (9)$$

A gauge-invariant two-point function describing the correlation of such a static-light meson is

$$G_\phi(x, y) = \langle \text{Tr} \phi^\dagger(t) U^{\text{fund}}(x, y) \phi(y) \rangle, \\ U^{\text{fund}}(x, y) = \mathcal{P} \exp(ig \int_x^y dx_i A_i^a T^a). \quad (10)$$

At large separations this two-point function falls off with the same mass parameter as in the Wilson loop perimeter law [23],

$$G_\phi(x, y) \sim e^{-M|x-y|}. \quad (11)$$

Considering instead the static potential in adjoint representation, string breaking occurs already in pure gauge theory, because the adjoint string can couple to gluons that may be pair produced. The corresponding final state corresponds to a bound state of the static source and some gluonic constituent and is described by

$$G_F(x, y) = \langle F_{ij}^a(x) U_{ab}^{\text{adj}}(x, y) F_{kl}^b(y) \rangle, \\ U_{ab}^{\text{adj}}(x, y) = 2 \text{Tr} T^a U^{\text{fund}}(x, y) T^b [U^{\text{fund}}(x, y)]^\dagger, \quad (12)$$

with now an adjoint Wilson line connecting the field strength tensors.

The correlators eqs. (10) and (12) are manifestly gauge-invariant because of the string inserted between the fields. If the string is chosen to be a straight line, then it can be gauged away to be unity in some axial gauge. In such a gauge and in a perturbative regime the correlators coincide with the propagators for the scalar and gauge fields. Does this connection also hold in a non-perturbative regime? If so, the masses of the dynamical fields should be related to the constituents discussed in the last section. However, there is one subtlety involved, in that the masses extracted from the correlators G_ϕ , G_F contain a logarithmic divergence due to the self-energy of the Wilson line representing the static source. Fortunately, due to superrenormalisability in three dimensions, the divergence can be computed perturbatively and subtracted [24,25], leaving us with a finite number that is only defined up to a finite additive renormalisation that has to be fixed by some prescription.

Fig.4 shows the mass values extracted from a lattice simulation of the F-F correlator in comparison with the 1^{--} bound state mass [25]. In the Higgs phase, the correlator indeed reproduces the mass of the W-boson as also obtained from the propagator or the gauge-invariant operator V , eq. (4), after its finite part has been fixed by matching it to the V -correlator at $y = -0.02$. In the symmetric phase it stays at a practically constant value of $\sim 0.75g_3^2$, which is about half the glueball mass $m_P = 1.60(4)g_3^2$ [9], and about twice the propagator mass $m_A = 0.46(3)g_3^2$ [15]. Since the field strength tensor contains two covariant derivatives, $F_{ik} = [D_i, D_k]$, this is again in good agreement with the naive constituent picture expectation and may be taken as another piece of evidence for a gluonic mass unit entering gauge-invariant quantities.

Finally, another manifestly gauge-invariant and physical quantity of three-dimensional gauge theories is the




FIG. 4. The lowest physical 1^{--} state (the W-boson in the Higgs phase) and the mass M'' extracted from $G_F(x, y)$.

string breaking scale r_b itself. Its size depends on the string tension and the mass of the dynamical particles that have to be produced to break the string. The breaking scale for the fundamental representation potential in the SU(2) Higgs model has been calculated in a recent lattice simulation [26], by extracting it from the turnover of the potential as shown in Fig. 5. The continuum extrapolation of those results gives

$$r_b g_3^2 \approx 8.5, \quad r_b^{-1} \approx 0.12 g_3^2. \quad (13)$$

Again for comparison, the lightest scalar bound state from Fig. 2 is $m_R = 0.839(15)g_3^2$, and the lightest glueball $m_P = 1.60(4)g_3^2$. A calculation of the adjoint representation potential in pure gauge theory is still in progress, but a fairly good estimate for its breaking scale may be obtained by comparing the final state energy extracted from the F-F correlator with the energy of the linear part in the potential extrapolated from Wilson loop calculations. The result [25] is of the same order of magnitude as that given in eq.(13).

In summary, pure gauge theory as well as the Higgs model contain non-perturbative, dynamical mass scales that are much smaller than any of the masses present in the spectrum. As of yet, it is not understood what is the origin for those small scales, but intriguing to speculate whether they may have something to do with the previously discussed propagator mass scales.

VII. THE DEBYE MASS

An important concept in the phenomenology of high temperature QCD is the static electric screening mass, or the Debye mass m_D . Although it has a leading order




FIG. 5. The energies of the ground state and the first excited state for the static potential in fundamental representation. The dashed lines indicate the location of twice the energy of the single meson state, as extracted from G_ϕ .

contribution that is perturbative [27], it couples to the three-dimensional magnetic sector in next-to-leading order, and hence requires a non-perturbative treatment as well. The Debye mass can be expanded as

$$m_D = m_D^{\text{LO}} + \frac{Ng_3^2}{4\pi} \ln \frac{m_D^{\text{LO}}}{g_3^2} + c_N g_3^2 + \mathcal{O}(g^3 T), \quad (14)$$

where $m_D^{\text{LO}} = (N/3 + N_f/6)^{1/2} gT$ and N_f is the number of flavours. The logarithmic part of the $\mathcal{O}(g^2)$ correction can be extracted perturbatively [27], but c_N and the higher terms are non-perturbative. To allow for a lattice determination, a non-perturbative definition was formulated in [24], employing the $SU(N)$ adjoint Higgs model as the dimensionally reduced effective theory. By integrating out the heavy adjoint Higgs this can be further reduced to the pure $SU(N)$ theory in three dimensions. The statement of [24] is that the coefficient c_N entering the Debye mass can be determined from the exponential fall-off of an adjoint Wilson line with appropriately chosen adjoint charge operators at the ends. The F-F correlator discussed in the last section is precisely such an operator. From its measurement in a lattice simulation of 3d pure $SU(2)$ in [25] one finds the complete non-perturbative $\mathcal{O}(g_3^2)$ corrections to the Debye mass with high precision to be $c_2 = 1.06(4)$. Judging from the precision of the result, the correlator G_F seems to be much more amenable to the analysis than other operators that have been studied in [28]. A calculation for $SU(3)$ is currently in progress.

VIII. CONCLUSIONS

Electric and magnetic screening properties of hot non-Abelian plasmas are non-perturbative phenomena and

closely related to the physics of confinement in three-dimensional gauge theories. The physical properties of $SU(2)$ pure gauge theory and the $SU(2)$ Higgs model in three dimensions have been studied by means of lattice simulations, and the mass spectrum as well as the static potential are known with rather good precision. The inverse string breaking scales of the potentials in adjoint and fundamental representation constitute physical mass scales which in both cases are much smaller than those of the lightest states in the spectrum. It was attempted to establish a connection between the magnetic mass, as obtained from simulations of the gluon propagator, and the mass spectrum as well as the static potential by means of a constituent picture. Although suggestive, the proposed connection is not conclusive at this level. In order to make progress in this direction, it would be necessary to find a gauge-invariant correlation function giving just one gluon constituent mass and thus reproducing the propagator simulations. Regarding the Debye mass, the full non-perturbative corrections of order $\sim g_3^2$ arising from three-dimensional pure gauge dynamics have been determined following the method of [24].

IX. ACKNOWLEDGMENTS

It is a pleasure to thank U. Heinz and the organisers for a stimulating conference.

-
- [1] P. Ginsparg, Nucl. Phys. B170 (1980) 388; T. Appelquist and R. Pisarski, Phys. Rev. D23 (1981) 2305.
 - [2] K. Rummukainen, Nucl.Phys.Proc.Suppl.53 (1997) 30.
 - [3] K. Kajantie, M. Laine, K. Rummukainen and M. Shaposhnikov, Nucl.Phys. B503 (1997) 357; A. Nieto, Int. J. Mod. Phys. A12 (1997) 1431.
 - [4] A. D. Linde, Phys. Lett. B96 (1980) 289; D. Gross, R. Pisarski and L. Yaffe, Rev. Mod. Phys. 53 (1981) 43
 - [5] K. Kajantie, M. Laine, K. Rummukainen and M. Shaposhnikov, Nucl. Phys. B458 (1996) 90.
 - [6] K. Rummukainen, M. Tsy-pin, K. Kajantie, M. Laine, M. Shaposhnikov, hep-lat/9805013. K. Rummukainen, these proceedings.
 - [7] W. Buchmüller and O. Philipsen, Nucl. Phys. B443 (1995) 47
 - [8] K. Kajantie, M. Laine, K. Rummukainen and M. Shaposhnikov, Phys. Rev. Lett. 77 (1996) 2887.
 - [9] O. Philipsen, M. Teper and H. Wittig, Nucl. Phys. B469 (1996) 445; Nucl. Phys. B528 (1998) 379.
 - [10] M. Teper, hep-lat/9804008.
 - [11] M. Teper, Phys. Lett. B311 (1993) 223.
 - [12] A. Hart and O. Philipsen, in preparation.
 - [13] V.P. Nair, these proceedings.

- [14] F. Karsch, T. Neuhaus, A. Patkós and J. Rank, Nucl. Phys. B474 (1996) 217.
- [15] F. Karsch, M. Oevers and P. Petreczky, hep-lat/9807035.
- [16] R. Kobes, G. Kunstatter and A. Rebhan, Nucl. Phys. B355 (1991) 1.
- [17] G. 't Hooft, proceedings of the 19th Schladming Winter School, p.531.
- [18] G. Alexanian and V.P. Nair, Phys. Lett. B352 (1995) 435.
- [19] R. Jackiw and S.Y. Pi, Phys. Lett. B403 (1997) 297.
- [20] J. Cornwall, Phys. Rev. D57 (1998) 3694.
- [21] F. Eberlein, hep-ph/9804460.
- [22] W. Buchmüller and O. Philipsen, Phys. Lett. B 397 (1997) 112.
- [23] H. Evertz et al., Phys. Lett. B194 (1987) 277.
- [24] P. Arnold and L. Yaffe, Phys. Rev. D52 (1995) 7208.
- [25] M. Laine and O. Philipsen, Nucl. Phys. B523 (1998) 267.
- [26] O. Philipsen and H. Wittig, hep-lat/9807020.
- [27] A.K. Rebhan, Phys. Rev. D48 (1993) R3967; Nucl. Phys. B430 (1994) 319.
- [28] K. Kajantie, M. Laine, J. Peisa, A. Rajantie, K. Rummukainen, M. Shaposhnikov, Phys. Rev. Lett. 79 (1997) 3130.